

continuously for 7 days or until signs of dysentery disappear.

(iii) *Limitations.* Prepare a fresh solution daily. Treatment not to exceed 14 days. If symptoms persist after 4 to 5 days consult a veterinarian. Not to be given to swine that weigh more than 250 pounds.

(4) *Growing quail*—(i) *Amount.* 400 milligrams per gallon in drinking water.

(ii) *Indications for use.* For prevention of ulcerative enteritis due to *Clostridium colinum* susceptible to bacitracin methylene disalicylate.

(iii) *Limitations.* Prepare fresh solution daily. Use as sole source of drinking water.

[57 FR 37322, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992, as amended at 63 FR 38474, July 17, 1998; 64 FR 13068, Mar. 17, 1999]

§ 520.154b Soluble bacitracin methylene disalicylate and streptomycin sulfate oral powder.

(a) *Specifications.* Each gram contains 200 units of soluble bacitracin methylene disalicylate, streptomycin sulfate equivalent to 20 milligrams of streptomycin, and 850 milligrams of carob flour.

(b) *Sponsor.* See No. 062925 in § 510.600(c) of this chapter.

(c) *Conditions of use.* *Dogs*—(1) *Amount.* 1 level teaspoonful per 10 pounds of body weight three times daily, mixed in a small quantity of liquid or feed.

(2) *Indications for use.* Treatment of bacterial enteritis caused by pathogens susceptible to bacitracin and streptomycin such as *Escherichia coli*, *Proteus* spp., *Staphylococcus* spp., and *Streptococcus* spp., and for the symptomatic treatment of associated diarrhea.

(3) *Limitations.* If no improvement is noted in 2 to 3 days, diagnosis should be reevaluated. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37322, Aug. 18, 1992, as amended at 61 FR 66581, Dec. 18, 1996]

§ 520.154c Bacitracin zinc soluble powder.

(a) *Specifications.* Each pound contains the equivalent of not less than 5 grams of bacitracin.

(b) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.70 of this chapter.

(d) *Conditions of use.* (1) *Broiler chickens*—(i) *Amount.* 100 milligrams per gallon in drinking water.

(A) *Indications for use.* Prevention of necrotic enteritis caused by *Clostridium perfringens* susceptible to bacitracin zinc.

(B) *Limitations.* Prepare a fresh solution daily.

(ii) *Amount.* 200 to 400 milligrams per gallon in drinking water.

(A) *Indications for use.* Control of necrotic enteritis caused by *Clostridium perfringens* susceptible to bacitracin zinc.

(B) *Limitations.* Prepare a fresh solution daily.

(2) *Growing quail*—(i) *Amount.* 500 milligrams per gallon in drinking water for 5 days followed by 165 milligrams per gallon in drinking water for 10 days.

(ii) *Indications for use.* Control of ulcerative enteritis caused by *Clostridium* spp. susceptible to bacitracin zinc.

(iii) *Limitations.* Prepare a fresh solution daily.

[57 FR 37322, Aug. 18, 1992, as amended at 67 FR 78355, Dec. 24, 2002]

§ 520.182 Bicyclohexylammonium fumagillin.

(a) *Specifications.* The drug is a soluble powder containing bicyclohexylammonium fumagillin and appropriate phosphate buffers.

(b) *Sponsor.* See No. 059620 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used for the prevention of nosema in honey bees.¹

(2) It is administered usually in a 2:1 sugar sirup containing a concentration of from 75 to 100 milligrams of fumagillin activity per gallon of sugar sirup.¹

(3) Colonies used for package production should be fed medicated sirup as a principal food supply for a month prior

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.